

Yaroslavl State University
Preprint YARU-HE-97/05
hep-ph/9709218

Radiative Transition of Massless Neutrino in Strong Magnetic Field

A.A. Gvozdev, N.V. Mikheev

Yaroslavl P.G. Demidov State University, Yaroslavl 150000, Russia
E-mail: gvozdev@univ.uniya.ac.ru mikheev@yars.free.net

and

L.A. Vassilevskaya

Moscow M.V. Lomonosov State University, V-952, Moscow 117234, Russia
E-mail: vasilevs@vitep5.itep.ru

Abstract

Radiative transition of massless neutrino $\nu_l \rightarrow \nu_l + \gamma$ ($l = e, \mu, \tau$) in a strong magnetic field is investigated in the framework of the Standard Model. The process probability and the mean values of the neutrino energy and momentum loss are presented. Possible astrophysical manifestation of the process considered is analysed.

*Talk given at the XXXII Rencontres de Moriond
on Electroweak interactions and unified theories,
Les Arcs, Savoie, France, 15-22 March 1997.*

The investigations of neutrino processes in strong magnetic fields are of principal interest in astrophysics where gigantic neutrino fluxes and strong magnetic fields can take place simultaneously (a process of a coalescence of neutron stars, an explosion of a supernova). By magnetic field is meant the field of strength of order and more the critical, so called, Schwinger value $B > B_s$, $B_s = m_e^2/e \simeq 0.44 \cdot 10^{14} G$. By now astrophysical knowledge of strong magnetic fields which can be realized in the nature have changed essentially. For example, the field strengths inside the astrophysical objects in principle could be as high as $10^{15} - 10^{18} G$ [1]. On the other hand, in such strong magnetic fields otherwise negligible processes are not only opened kinematically but become substantial ones as well. For example, a photon splitting into two photons $\gamma \rightarrow \gamma + \gamma$ [2], a magnetic catalysis of the massive neutrino radiative decay [3].

The diagonal radiative neutrino transition becomes possible due to the photon dispersion in the strong magnetic field which differs significantly from the vacuum dispersion with increasing photon energy. So the real photon 4-momentum can appear as a space-like and sufficiently large one ($|q^2| \gg m_\nu^2$). In this case the phase space for diagonal neutrino transition $\nu_l \rightarrow \nu_l + \gamma$ ($l = e, \mu, \tau$) is opened also. Thus this diagonal radiative neutrino transition manifests itself as neutrino bremsstrahlung, does not contain uncertainties associated with a possible mixing in the lepton sector of the SM and can lead to observable physical effects in the strong magnetic fields $B \gg B_s$.

The important property of the amplitude $\nu \rightarrow \nu + \gamma$ we have studied is that it has a square-root singularities of the type:

$$\sim (\mathcal{E}_{nm}^2 - q_{\parallel}^2)^{-1/2},$$

in the threshold points $(q_{\parallel}^2)_{thr} = \mathcal{E}_{nm}^2$, labelled by n, m ($n, m = 0, 1, 2, \dots$):

$$\mathcal{E}_{nm}^2 = \left(\sqrt{m_e^2 + 2eBn} + \sqrt{m_e^2 + 2eBm} \right)^2,$$

where $q_{\parallel}^2 = q_0^2 - q_3^2$ in the frame with $\vec{B} = (0, 0, B)$.

$\mathcal{E}_{nm}$ has the simple physical meaning as transversal energy of an electron-positron pair in the magnetic field. The similar phenomena is known as the cyclotronic resonance in the vacuum polarization [4].

It is of interest for some astrophysical applications the case of relatively high neutrino energy $E_\nu \simeq 10 - 20 \text{ MeV} \gg m_e$ and strong magnetic field $eB \gg E_\nu^2$, when a region of the cyclotronic resonance on the ground Landau level $q_\parallel^2 \rightarrow 4m_e^2$ dominates in this process. The square-root singularities on higher Landau levels are not realized under the condition $eB > E_\nu^2$.

The main contribution to the probability of high energy neutrino ($eB \gg E_\nu^2 \gg 4m_e^2$) is determined from the vicinity of the resonance point $q_\parallel^2 = 4m_e^2$ and can be presented in the form:

$$W \simeq \frac{\alpha G_F^2}{8\pi^2} (g_v^2 + g_a^2) (eB)^2 E_\nu \sin^2 \theta, \quad (1)$$

where θ is the angle between the vectors of the magnetic field strength $\vec{B}$ and the momentum of the initial neutrino $\vec{p}$; g_v and g_a are the well-known vector and axial-vector $\nu - e$ coupling constants in the SM.

It is of more practical importance for some astrophysical applications to calculate the four-vector of the neutrino energy and momentum loss in the process of the neutrino radiative transition:

$$Q_\mu = E_\nu \int dW \cdot q_\mu. \quad (2)$$

The physical meaning of this vector is that Q_0/E_ν is the mean energy loss in a unit time and $\vec{Q}/E_\nu$ is the momentum loss in a unit time. We have obtained the expression for the vector Q_μ :

$$Q_\mu = \frac{1}{4} E_\nu W [p_\mu - (p\varphi\varphi)_\mu - \frac{2g_v g_a}{g_v^2 + g_a^2} (p\tilde{\varphi})_\mu], \quad (3)$$

where $\varphi_{\alpha\beta} = F_{\alpha\beta}/B$ and $\tilde{\varphi}_{\alpha\beta} = \frac{1}{2}\varepsilon_{\alpha\beta\mu\nu}\varphi_{\mu\nu}$ are the dimensionless tensor and dual tensor of the magnetic field.

To illustrate the applications of the results we have obtained we estimate below possible effects of this process in astrophysical cataclysm of type of a supernova explosion or a merger of neutron stars under the condition that a compact remnant has for some reasons extremely strong magnetic field, $B \simeq 10^{16} - 10^{18} G$. We assume that neutrinos of all species with the typical mean energy $\bar{E}_\nu \simeq 20 \text{ MeV}$ are radiated from a surface of a neutrinosphere in such a process. Using (3) we estimate the neutrino energy loss:

$$\mathcal{E} \simeq 10^{50} \left(\frac{\mathcal{E}_{tot}}{10^{53} erg} \right) \left(\frac{B}{10^{17} G} \right)^2 \left(\frac{\bar{E}_\nu}{20 MeV} \right) \left(\frac{R}{10 km} \right) erg,$$

where $\mathcal{E}_{tot} \simeq 10^{53}$ erg is the total neutrino radiation energy; B is the magnetic field strength in the vicinity of the neutrinosphere of radius R . Neutrino bremsstrahlung γ -quanta are captured by a strong magnetic field and propagate along the field [4]. Thus the mechanism of significant power "pumping" of polar caps of a magnetized remnant can take place. This phenomenon can result in reemission within a narrow cone $\Omega/4\pi \ll 1$ along the magnetic moment of the remnant. So, if an external absorbing envelope is absent in such an extraordinary astrophysical cataclysm (very strong magnetic fields, gigantic neutrino fluxes) the reemission process will be an observable effect. Namely, such a reemission in the magnetic field could appear as an anisotropic γ -burst with the duration of order of the neutrino emission time and of the typical energy $\mathcal{E}_{4\pi}^{burst} \sim 10^{51}$ erg in terms of 4π -geometry.

The work of N.V. Mikheev was supported by a Grant N d104 from the International Soros Science Education Program. The work of L.A. Vassilevskaya was supported by a fellowship of INTAS Grant 93-2492-ext and was carried out within the research program of International Center for Fundamental Physics in Moscow.

References

- [1] M. Ruderman, in *Neutron Stars: Theory and Observation*, ed. by J. Ventura and D. Pines, Kluwer Academic. Pub., Dordrecht, 1991;
V.V. Usov, Nature 357 (1992) 472;
M. Bocquet et al., Astr. and Ap. 301 (1995) 757.
- [2] S.L. Adler, Ann. Phys. N.Y. 67 (1971) 599.
- [3] A.A. Gvozdev, N.V. Mikheev and L.A. Vassilevskaya, Phys. Rev. D54 (1996) 5674.
- [4] A.E. Shabad, Ann. Phys. N.Y. 90 (1975) 166;
A.E. Shabad and V.V. Usov, Astrophysics and Space Science, 102 (1984) 327.